

Cosmological Constant and Supergravity Soft Terms*

Jihn E. Kim

Department of Physics and Center for Theoretical Physics, Seoul National University

Seoul 151-742, Korea

Abstract

In supergravity models, the quantum correction to the vacuum energy can be of order M_I^4 , if the cutoff is of order the Planck mass M_P^2 and $Str \mathcal{M}^2 \neq 0$. Therefore, the tree level cosmological constant must be nonzero (probably negative). Since the current practice of calculating soft parameters in supergravity models assumes the vanishing tree level cosmological constant, the supergravity calculation must be accordingly modified. This implies that the soft parameters in supergravity has an additional contribution depending on the nonvanishing tree level cosmological constant. A simple quantum mechanical model mimicking this situation is also presented.

Typeset using REVTeX

*Talk presented at the ITP Workshop on “Frontiers in QFT”, Beijing, June 15–24, 1994.

There are three well-known hierarchy problems: the cosmological constant problem, the Kaluza–Klein problem, and the gauge hierarchy problem. If the theoretical physics in 1917 were as advanced as now, the introduction of the cosmological constant [1] might have had led to the immediate theoretical problem, the cosmological constant problem. In the same vein, the Kaluza–Klein theory [2] introduced a huge ratio, $M_P/m_e \gg 1$, working against the original beautiful idea, which is the reason that the Kaluza–Klein theory seems currently to be in disfavor. The third problem is the well-known gauge hierarchy problem in grand unified theories [3]. In this talk I discuss the cosmological constant in supergravity, and hope that it instigate young Chinese physicists to think about this eighty years old problem again and get an idea toward the solution. Certainly my talk is not dealing with a solution, but can give an idea toward a correct direction of the solution.

This talk is based on my recent work collaborated with Kiwoon Choi and Hans Peter Nilles on the cosmological constant in supergravity and its effect to soft terms [4].

Supersymmetry (SUSY) is widely believed to be a leading candidate for a further symmetry beyond the standard model. This is largely due to the fact that SUSY provides the only known perturbative solution to the problem of quadratic divergence in the Higgs boson mass. The most popular working scenario is supergravity models in which SUSY is broken at the intermediate scale [5].

At low energy, the effective theory is parametrized by SUSY breaking soft parameters in addition to the renormalizable couplings allowed by the gauge symmetry. An interesting feature of supergravity models is that many of the coefficients of soft terms are calculable at tree approximation. In some cases, one obtains certain tree level relations among the soft coefficients renormalized at the Planck scale M_P . For instance, one obtains $A = -\sqrt{3}m_0\lambda$ in the dilaton-dominated SUSY breaking scenario in string theory [6, 7], where A denotes a generic trilinear scalar coefficient, λ is the associated Yukawa coupling, and m_0 is the soft scalar mass.

For physical applications of such tree level results, one has to take into account quantum corrections. Ordinary renormalizable interactions lead to one loop corrections proportional to

$$\frac{g^2}{8\pi^2} \ln(\Lambda^2/\mu^2) \quad (1)$$

where g denotes a gauge or Yukawa coupling constant, μ is a scale around the weak scale m_W where the loop graph is being evaluated, and Λ is the momentum cutoff above which the validity of four-dimensional $N = 1$ supergravity theory breaks down. Clearly for g^2 of order unity and Λ not far below M_P , the corrections become important due to a large logarithm. A nice feature of this type of corrections is that they depend on the cutoff Λ logarithmically and are *calculable* within the supergravity model by the aid of renormalization group analysis.

Besides the above type of calculable corrections, there are other types of corrections which depend strongly on Λ and thus whose precise magnitudes are *not* calculable within the supergravity model [8]. For instance, one loop graphs induced by nonrenormalizable gravitational interactions would give power-law divergent corrections which are proportional to

$$\frac{1}{8\pi^2} (\kappa\Lambda)^n, \quad (2)$$

where $\kappa = \sqrt{8\pi}/M_P$ denotes the dimensionful gravitational coupling constant and n is a positive integer. Obviously, the main correction in this case arises from the fluctuations at the scale Λ . As a result, they are strongly dependent on the unknown physics at the scale Λ . Thus one has to know the underlying theory in more detail to get a useful information on such parameters.

Roughly speaking, the dimensionless coupling ‘ $\kappa\Lambda$ ’ measures how strongly the underlying theory of supergravity is coupled. If strongly coupled, one can guess that $\kappa\Lambda \simeq 4\pi$ which means that loops are as important as trees. (This is analogous to the nonlinear sigma model of the pions whose underlying theory is the strongly coupled QCD [9].) In this case, tree level predictions would not be useful at all *unless* they are mere consequence of some symmetries of the underlying theory.

Of course, one can ignore the incalculable power-law divergent corrections if the underlying theory is very weakly coupled, i.e. $\kappa\Lambda \ll 1$. However, a more reasonable scenario would be that the underlying theory is moderately weakly coupled with $\kappa\Lambda$ of order unity, giving a loop factor of order $1/8\pi^2$. This is indeed the case for supergravity models which correspond to the low energy limit of string theory. In this case, it is natural to set Λ to the string scale ¹ M_{string} . In the heterotic string theory, the gravitational coupling constant κ is given by $\kappa M_{\text{string}} = g_{\text{GUT}} = O(1)$ [10] where g_{GUT} is the unified gauge coupling constant at M_{string} . In view of the string theory, thus the interesting case is a moderately weakly coupled supergravity, leading to $\kappa\Lambda$ being of order unity.

For $\kappa\Lambda = g_{\text{GUT}}$, the power-law divergent corrections of Eq. (2) are expected to be of order $\alpha_{\text{GUT}}/2\pi$ and thus are not so significant. This seems to be a small correction. But if there are sufficiently large number of particles in the loop, then the correction can be of $O(1)$. Without supersymmetry, however, it is difficult to take this effect into account.

With softly broken supersymmetry, the radiative correction to the coefficients of operators $\mathbf{1}$, $\phi^*\phi$ and $\bar{\psi}\psi\phi$ are quadratically divergent, logarithmically divergent and finite, respectively. The coefficient of $\mathbf{1}$ is the cosmological constant. The cosmological constant is an important unresolved problem in spontaneously broken gauge theories [11]. At one loop order, the cosmological constant which is the vacuum energy density in our notation receives a quadratically-divergent zero point energy contribution [12]. For N chiral multiplets with $m_B \gg m_F$, the contribution is given by

$$N \int \frac{d^3k}{(2\pi)^3} \left(\sqrt{\mathbf{k}^2 + m_B^2} - \sqrt{\mathbf{k}^2 + m_F^2} \right) \simeq N \frac{m_B^2 \Lambda^2}{8\pi^2}. \quad (3)$$

This leads to corrections of order $N\alpha_{\text{GUT}}/2\pi$ to some of soft coefficients. A key point in this regard is that although $\alpha_{\text{GUT}}/2\pi$ is small, $N\alpha_{\text{GUT}}/2\pi$ can be significantly large since in realistic models N is typically of order $8\pi^2$. ² Therefore, the cosmological constant is shifted by an amount of order M_I^4 which is a natural order of the tree level cosmological constant. Therefore, calculations of soft parameters assuming a vanishing tree level cosmological constant can be significantly affected.

To be definite, let us consider a simple supergravity model with the following Kähler potential and superpotential [5]

$$\begin{aligned} K &= h_\alpha h_\alpha^* + \phi_i \phi_i^* + (\xi H_1 H_2 + h.c.) , \\ W &= W_h(h_\alpha) + \frac{1}{6} \tilde{\lambda}_{ijk} \phi_i \phi_j \phi_k + \tilde{\mu} H_1 H_2, \end{aligned} \quad (4)$$

¹It has been argued [10] that another candidate for the cutoff, the compactification scale, is quite close to M_{string} .

²In the minimal supersymmetric standard model $N = 49$, but it can be larger if there are more chiral multiplets at the intermediate scale.

where h_α denote hidden sector fields triggering SUSY breaking, and ϕ_i are generic observable sector fields including quarks, leptons, and the two Higgs doublets H_1 and H_2 .

To obtain the effective action S_ϕ of the observable fields ϕ_i , we integrate out the hidden sector fields to obtain

$$\exp(iS_\phi) = \int [\mathcal{D}h] \exp(iS), \quad (5)$$

where S is the full supergravity action, and $[\mathcal{D}h]$ includes the integration of the gravity multiplet $(g_{\mu\nu}, \psi_\mu)$ over a background spacetime metric $\bar{g}_{\mu\nu}$ with macroscopic wavelength. Since the high momentum modes of ϕ_i are not integrated out yet, S_ϕ is defined at the cutoff scale Λ in the sense of Wilson. Thus to study the low energy physics of ϕ_i , one still needs to scale the renormalization point down to the weak scale. At any rate, in the flat limit, S_ϕ would be characterized by the effective superpotential of global SUSY [5]

$$W_{\text{eff}} = \frac{1}{6} \lambda_{ijk} \phi_i \phi_j \phi_k + \mu H_1 H_2, \quad (6)$$

and also the soft breaking part of the form

$$\mathcal{L}_{\text{soft}} = m_0^2 \phi_i \phi_i^* + \left(\frac{1}{6} A_{ijk} \phi_i \phi_j \phi_k + B H_1 H_2 + \text{h.c.} \right). \quad (7)$$

Among the coefficients in W_{eff} and $\mathcal{L}_{\text{soft}}$, those which depend on the expectation value of the hidden sector scalar potential are relevant for us. They are the soft scalar mass m_0^2 and the B coefficient. To calculate these, we expand the supergravity potential V in powers of light fields $\phi^* \phi$. Noting that

$$V = e^G \left[\frac{\partial G}{\partial z_I} (G^{-1})^J_I \frac{\partial G}{\partial z_J^*} - 3 \right] \quad (8)$$

where

$$G = K + \log |W|^2, \quad (9)$$

we obtain

$$\begin{aligned} V = & V_h + (m_{3/2}^2 + \kappa_0^2 V_h) \phi_i^* \phi_i \\ & + [(1 + |\xi|^2) m_{3/2}^2 + \kappa_0^2 V_h] (H_1^* H_1 + H_2^* H_2) \\ & + (2m_{3/2}^2 + \kappa_0^2 V_h) (\xi H_1 H_2 + \xi^* H_1^* H_2^*) + O(\phi_i^3). \end{aligned} \quad (10)$$

By taking the expectation values of the coefficients of light fields, the soft parameters are given by

$$\begin{aligned} m_0^2 &= \langle m_{3/2}^2 + \kappa_0^2 V_h \rangle \\ m_H^2 &= \langle (1 + |\xi|^2) m_{3/2}^2 + \kappa_0^2 V_h \rangle \\ B &= \langle \xi (\kappa_0^2 V_h + 2m_{3/2}^2) \rangle. \end{aligned} \quad (11)$$

The other soft parameters and couplings are given by

$$\begin{aligned} \lambda_{ijk} &= \langle \tilde{\lambda}_{ijk} \exp(\kappa_0^2 h_\alpha h_\alpha^* / 2) \rangle, \\ \mu &= \langle (\tilde{\mu} + \xi \kappa_0^2 W_h) \exp(\kappa_0^2 h_\alpha h_\alpha^* / 2) \rangle, \\ A_{ijk} &= \langle \tilde{\lambda}_{ijk} \kappa_0^2 (h_\alpha D_\alpha W_1)^* \exp(\kappa_0^2 h_\alpha h_\alpha^*) \rangle, \end{aligned} \quad (12)$$

where κ_0 is the bare gravitational coupling constant, $D_\alpha W_h = (\partial_{h_\alpha} + \kappa_0^2 h_\alpha^*) W_h$, and the gravitino mass $m_{3/2}$ and the hidden sector scalar potential V_h are given by

$$\begin{aligned} m_{3/2}^2 &= \kappa_0^4 |W_h|^2 \exp(\kappa_0^2 h_\alpha h_\alpha^*), \\ V_h &= (|D_\alpha W_h|^2 - 3\kappa_0^2 |W_h|^2) \exp(\kappa_0^2 h_\alpha h_\alpha^*). \end{aligned} \quad (13)$$

Here the bracket means the average over the hidden sector fields. For example,

$$\langle V_h \rangle = \int [\mathcal{D}h] V_h(h_\alpha) \exp(iS_h) / \int [\mathcal{D}h] \exp(iS_h), \quad (14)$$

where S_h is the supergravity action of the hidden sector fields alone. If interactions among hidden sector fields are weak enough, which is usually the case,³ the above discussion applies. The soft parameters m_0^2 and B depends on $\langle V_h \rangle$. But $\langle V_h \rangle$ is not the cosmological constant.

Then, what is the cosmological constant? The fully renormalized cosmological constant V_{eff} at low energy is obtained by integrating out all the fields in the theory,⁴

$$\exp(i \int d^4x V_{\text{eff}}) = \int [\mathcal{D}\phi \mathcal{D}h] \exp(i \int d^4x \sqrt{g} \mathcal{L}) \quad (15)$$

where $[\mathcal{D}\phi]$ represents the integration over all the observable gauge and matter multiplets.

Of course, $\langle V_h \rangle$ and V_{eff} are different. At tree level, however, they are the same. In the classical approximation, $\langle V_h \rangle$ is simply the classical potential in $\mathcal{L}_h$ since the path is taken over the fields satisfying the classical equation of motion. From Eq. (15), the tree level value of V_{eff} is just the classical potential in $\mathcal{L}_h$ since the nontrivial contribution of $[\mathcal{D}\phi]$ integration is the effects of loops of ϕ fields. Therefore, at tree level the cosmological constant V_{eff} and $\langle V_h \rangle$ are the same.

Note, however, that although $\langle V_h \rangle_{\text{tree}}$ corresponds to the *tree level* cosmological constant, the quantity $\langle V_h \rangle$ that appears in the soft coefficients of Eq. (11) is *not* the fully renormalized cosmological constant. Eq. (11) shows that m_0^2 and the part of B associated with the Kähler potential term ($\xi H_1 H_2 + h.c.$) [13] depend on $\langle V_h \rangle$. For local SUSY broken by the vacuum value of the auxiliary component $F_\alpha = D_\alpha W_h \exp(\kappa_0^2 h_\alpha h_\alpha^*/2)$, *unless* one implements fine tuning of some parameters in the hidden sector superpotential W_h , the typical size of $\langle V_h \rangle$ would be of $O(\langle |F_\alpha|^2 \rangle) = O(\kappa_0^{-2} m_{3/2}^2)$. In most of the previous studies, motivated by the fully renormalized vanishing cosmological constant, the tree level expectation value of the hidden sector scalar potential, $V_0 \equiv \langle V_h \rangle_{\text{tree}}$, was assumed

³ In some cases, hidden sector contains gauge interactions which become strong to provide a nonperturbative seed, e.g. the gaugino condensation [15], for SUSY breaking. We assume in such cases that the strongly interacting gauge nonsinglet sector is already integrated out, whose effects are included in the effective hidden sector superpotential W_h of weakly interacting gauge singlet h_α .

⁴The long wave length metric $\bar{g}_{\mu\nu}$ is treated as background and is not integrated out to see the gravitational effect at low energy.

to be zero or at least $\kappa_0^2 V_0 \ll m_{3/2}^2$. However, as we have already anticipated in Eq. (3), for a large number of chiral multiplets, $N = O(8\pi^2)$, and the choice of the cutoff, $\kappa_0 \Lambda = O(1)$, quadratically divergent quantum correction to the vacuum energy density becomes of order $\kappa_0^{-2} m_{3/2}^2$. This implies $\kappa_0^2 V_0 = O(m_{3/2}^2)$, and then one can not ignore⁵ the $\langle V_h \rangle$ -dependent part of m_0^2 and B in Eq. (11).

At one loop, the vacuum energy density receives a contribution of the form [12]

$$V_{\text{eff}} - (V_{\text{eff}})_{\text{tree}} = \int \frac{d^4 p}{(2\pi)^4} \text{Str} [\ln(p^2 + \mathcal{M}^2)] \simeq \frac{1}{8\pi^2} \text{Str} (\mathcal{M}^2) \Lambda^2. \quad (16)$$

Here $(V_{\text{eff}})_{\text{tree}}$ is the tree level value of V_{eff} , thus it equals $\langle V_h \rangle_{\text{tree}}$. For our simple model, we have

$$\text{Str} (\mathcal{M}^2) \simeq (Nm_0^2 - \tilde{N}\tilde{m}^2) + (\text{hidden sector contribution}) \equiv N_{\text{eff}} m_0^2, \quad (17)$$

where N is the number of observable chiral multiplets, $\tilde{N}$ is the number of gauginos which are assumed to have a common mass $\tilde{m}$, and we neglected the masses of matter fermions and gauge bosons. For the contribution from hidden sector, the gravity sector gives a negative contribution ($= -4m_{3/2}^2$) while hidden chiral multiplets give a positive contribution proportional to the number of hidden multiplets. Using Eq. (11), one easily finds

$$V_{\text{eff}} - (V_{\text{eff}})_{\text{tree}} \simeq \frac{N_{\text{eff}}}{8\pi^2} m_0^2 \Lambda^2 \simeq \frac{N_{\text{eff}}}{8\pi^2} \Lambda^2 (m_{3/2}^2 + \kappa_0^2 \langle V_0 \rangle). \quad (18)$$

Then the requirement of $V_{\text{eff}} = 0$ leads to

$$\kappa_0^2 \langle V_h \rangle_{\text{tree}} \simeq -\frac{\epsilon}{1 + \epsilon} m_{3/2}^2, \quad (19)$$

where

$$\epsilon = \frac{N_{\text{eff}}}{8\pi^2} (\kappa_0 \Lambda)^2. \quad (20)$$

Unlike the cosmological constant, $\langle V_h \rangle$ receives contributions only from the hidden field fluctuations. For $\kappa_0 \Lambda = g_{\text{GUT}}$, if (i) interactions among hidden sector fields are weak and (ii) the number of hidden multiplets which contribute to supersymmetry breaking is $O(1)$, we have

$$\delta \langle V_h \rangle = O\left(\frac{\alpha_{\text{GUT}}}{\pi} |F_\alpha|^2\right) = O\left(\frac{\alpha_{\text{GUT}}}{\pi} \kappa_0^{-2} m_{3/2}^2\right) \quad (21)$$

where $\delta \langle V_h \rangle = \langle V_h \rangle - \langle V_h \rangle_{\text{tree}}$, and F_α is the F -term of h_α , $F_\alpha = D_\alpha W_h \exp(\kappa_0^2 h_\beta h_\beta^*/2) = O(\kappa_0^{-2} m_{3/2}^2)$. Note that conditions (i) and (ii) are satisfied by many simple hidden sector models.

⁵ Recently Brignole, Ibanez and Munoz [7] also discussed the dependence of soft parameters on $\langle \kappa_0^2 V_h \rangle$ in the context of string-inspired supergravity models. However, they did not provide any rationale for $\langle \kappa_0^2 V_h \rangle$ to be of order $m_{3/2}^2$.

Thus we expect that $V_{\text{eff}} \neq \langle V_h \rangle$ and the difference is $O(\kappa_0^{-2} m_{3/2}^2)$.

Since the gaugino mass contribution is expected to be significantly smaller than the chiral matter contribution, it is quite conceivable that N_{eff} is *positive* and of $O(8\pi^2)$. Note that N_{eff} receives a contribution from all chiral multiplets with masses far below M_P , particularly from those in the minimal supersymmetric standard model with 49 chiral multiplets. Then for $\kappa_0 \Lambda = g_{\text{GUT}}$, which is motivated by string theory, ϵ is essentially of order unity. This implies that the soft scalar mass m_0^2 and the part of B associated with the Kähler potential term $(\xi H_1 H_2 + \text{h.c.})$ can be significantly affected by the contribution from $\kappa_0^2 \langle V_h \rangle$. This also means that to have a fully renormalized vanishing cosmological constant, the hidden sector scalar potential V_h is required to have a *negative* expectation value $\langle V_h \rangle$ of $O(\kappa_0^{-2} m_{3/2}^2)$ (see Eq. (19)). In this regard, we note that, in the ‘racetrack’ model [16] of gaugino condensations for SUSY breaking in string theory, the dilaton potential appears often to have a negative minimum value. Our discussion here indicates that a negative minimum of the dilaton potential is not a problem, but is a rather desirable feature for the fully renormalized cosmological constant to vanish. In passing, we note that the cosmological constant cannot be solved purely from the high energy physics alone; one has to deal with the quadratic divergence of the minimal supersymmetric standard model. If the tree level cosmological constant is zero, then the full cosmological constant is not zero.

One can mimick the above phenomenon in a simple quantum mechanical toy model. Let us introduce $(N + 1)$ dynamical variables, $\{p_0, q_0\}$ and $\{p_i, q_i\}$ ($i = 1, \dots, N$),

$$H = \frac{1}{2} p_0^2 + (E_0 + \frac{1}{2} \omega_0^2 q_0^2) + \sum_{i=1}^N \left[\frac{1}{2} p_i^2 + \frac{1}{2} \left\{ \omega^2 + \lambda \omega_0 (E_0 + \frac{\omega_0^2 q_0^2}{2}) \right\} q_i^2 \right] \quad (22)$$

where E_0, ω_0, ω and λ are parameters. This toy model is written such that the following correspondence between the previous supergravity model and the toy model makes a sense:

Hidden sector potential energy operator, $V_h \equiv E_0 + \frac{1}{2} \omega_0^2 q_0^2$

Mass (or frequency) of observable variables, $m^2 \equiv \omega^2 + \lambda \omega_0 \langle V_h \rangle$

The hidden sector ground state energy, $\tilde{V} \equiv \langle \frac{1}{2} p_0^2 + V_h \rangle$

The true ground state energy, $V_{\text{eff}} \leftrightarrow \langle H \rangle$

In the toy model, the classical ground state energy is given by the condition, $p_0 = p_i = q_0 = q_i = 0$. Namely, the tree level values are

$$\langle V_h \rangle_{\text{tree}} = E_0, \quad (V_{\text{eff}})_{\text{tree}} = E_0. \quad (23)$$

In this toy model also, the tree level values of $\langle V_h \rangle$ and V_{eff} are the same. As before, we want to satisfy $V_{\text{eff}} = 0$, not $(V_{\text{eff}})_{\text{tree}} = 0$.

If we require $\langle V_h \rangle_{\text{tree}} = (V_{\text{eff}})_{\text{tree}} = 0$, then

$$(m^2)_{\text{tree}} = \omega^2. \quad (24)$$

Now we can include the quantum effects. For ease of demonstration, let us assume

$$\omega_0 \simeq \omega, \quad \lambda \simeq \frac{1}{N} \simeq \frac{\omega}{E_0} \simeq \frac{1}{8\pi^2}. \quad (25)$$

Then

$$\langle H_h \rangle = \langle \frac{1}{2} p_0^2 + E_0 + \frac{1}{2} \omega_0^2 q_0^2 \rangle = \langle E_0 + \frac{\omega_0}{2} \rangle = E_0 \left(1 + O\left(\frac{1}{8\pi^2}\right) \right) \quad (26)$$

and

$$\langle V_h \rangle = \langle E_0 + \frac{1}{2}\omega_0^2 q_0^2 \rangle = \langle E_0 + \frac{\omega_0}{4} \rangle = E_0 \left(1 + O\left(\frac{1}{8\pi^2}\right) \right). \quad (27)$$

After performing the Gaussian integration with $\{p_0, q_0\}$, we obtain the low energy Hamiltonian,

$$H_\phi = \left(\sum_{i=1}^N \frac{1}{2} p_i^2 \right) + \left[E_0 + \frac{1}{2}\omega_0 + \frac{1}{2}(\omega^2 + \lambda\omega_0 E_0) \sum_{i=1}^N q_i^2 \right] + O(\lambda^2) \quad (28)$$

from which we can read the effective potential

$$V_{\text{eff}} = E_0 + \frac{1}{2}N\sqrt{\omega^2 + \lambda\omega_0 E_0} + O(\omega). \quad (29)$$

Thus, we obtain

$$V_{\text{eff}} - \langle V_h \rangle = \frac{1}{2}N\sqrt{\omega^2 + \lambda\omega_0 E_0} + O(\omega) \quad (30)$$

which has our previous form of $(1/8\pi^2) \times (\text{number of chiral fields})$.

In the above discussion, we considered only the one loop contribution (Fig. 1) to the vacuum energy density, which is of order $\kappa_0^{-2} m_{3/2}^2$ due to a large value of N_{eff} compensating over the loop suppression factor $1/8\pi^2$. Clearly this contribution persists even when all interactions (of course except for the kinetic and mass part) in the supergravity action are turned off. Inclusion of interactions gives rise to additional quadratic divergences, but at higher loop order. Contrary to the one loop contribution, these higher loop effects do *not* contain any additional large factor which may compensate over the additional loop suppression factor. Obviously two loop diagrams involving gauge interactions give a contribution smaller than the one loop effect by the small factor $\alpha_{\text{GUT}}/2\pi$. Note that quadratically divergent effects are dominated by the contribution from fluctuations at the cutoff scale where all interactions are presumed to be perturbative.

Fig. 1. One loop contribution to the cosmological constant.

One can consider other types of two loop diagrams involving trilinear and/or Yukawa scalar interactions. (See Figs. 2 and 3.) The contribution of Fig. 2 is roughly

$$\frac{1}{(4\pi^2)^2} \sum_{ijk} A_{ijk} A_{ijk}^* \Lambda^2, \quad (31)$$

while that of Fig. 3 is

$$\frac{1}{(4\pi^2)^2} \sum_{ijk} \lambda_{ijk} \lambda_{ijk}^* m_{3/2}^2 \Lambda^2. \quad (32)$$

Usually A_{ijk} is of order $m_{3/2} \lambda_{ijk}$, and then clearly the above two loop effects are negligible compared to the one loop effect of Eq. (18).

Fig. 2. The A term contribution at two loop level.

Fig. 3. Two loop contribution from Yukawa interaction.

Since the quantum corrections to the cosmological constant of order $\kappa_0^{-2} m_{3/2}^2 \gg m_W^4$ are added to the tree level value, one might wonder whether the vacuum structure of the observable sector fields ϕ_i is changed. Our results of Eqs. (11) and (18) explicitly show that soft masses of ϕ_i at the cutoff scale are still *positive* even after including the contribution from a *negative* $\langle V_h \rangle$. It is also easy to see that $\text{Str}(\mathcal{M}^2)$ is *independent* of ϕ_i and thus the vacuum structure of ϕ_i is untouched by the quadratically divergent corrections discussed above. To see it more explicitly, we can consider a simple example with the superpotential $W = \lambda \phi^3 + m \phi^2$ and the soft term $\mathcal{L}_{\text{soft}} = m_0^2 \phi \phi^*$. We then have,

for an arbitrary value of ϕ ,

$$\text{Str}(\mathcal{M}^2) = m_0^2 + |6\lambda\phi + 2m|^2 - |6\lambda\phi + 2m|^2,$$

which is ϕ -independent. Clearly this is a simple consequence of the nonrenormalization theorem for the effective potential in the presence of softly broken SUSY, particularly the absence of a field-dependent quadratic divergence.

So far we have shown that the quantum corrections to the cosmological constant in supergravity models can be of order $\kappa_0^{-2}m_{3/2}^2$ and most likely will be positive. This is for the reasonable choice of the cutoff scale, $\kappa_0\Lambda = O(g_{\text{GUT}})$, and mainly due to a large number of chiral multiplets which compensates over the loop suppression factor. This implies that conventional studies of supergravity phenomenology based on the input $\kappa_0^2\langle V_h \rangle \ll m_{3/2}^2$ should be modified. This also implies that it is rather desirable, for the fully renormalized vanishing cosmological constant to vanish, to have a negative tree level cosmological constant of order $\kappa_0^{-2}m_{3/2}^2$.

In our scheme, the squark masses m_i can be expressed as

$$\frac{m_i^2}{m_{3/2}^2} = 3C^2 \left\{ 1 + \frac{1}{3}N_i(T, T^*) \cos^2 \theta \right\} - 2 \quad (33)$$

where $\cos \theta$ signifies the relative importance in supersymmetry breaking by moduli and dilaton superfields (e.g. $\cos \theta \sim 0$ corresponds to the dilaton dominated supersymmetry breaking and vice versa) and N_i is related to the curvature of the Kähler manifold. For (2,2) Calabi-Yau spaces, $N_i \rightarrow -1$ as the moduli field $T \rightarrow \infty$. For orbifolds, N_i corresponds to modular weights of charged fields which are normally negative integer numbers [18]. Not to break $SU(3)_c \times U(1)_{\text{em}}$ gauge symmetry, we must require

$$C^2 > \frac{2}{3 + N_i(T, T^*) \cos^2 \theta}. \quad (34)$$

Because $\langle V_h \rangle$ most probably will be negative as discussed in Eq. (29), we would have $C^2 < 1$ (viz. $C^2 = 1 + (8\pi\langle V_h \rangle / 3m_{3/2}^2 M_{\text{Pl}}^2)$). At present, it is difficult to give any precise value for C^2 in view of the lack of information on the mass spectra of superpartners, and hence the phenomenological considerations studied for the case $C^2 = 1$ should serve only as a guideline [19]. For scalar masses, this contribution can shift them by $O(10)\%$, depending on the magnitude of ϵ .

In conclusion, I discussed the quantum correction to the cosmological constant in supergravity and its effects on the soft parameters. Even though the $N = 1$ supergravity is not renormalizable, this calculation seems to be reliable if the theory is replaced by a well behaved one above the cutoff scale. Without supersymmetry, this loop calculation has the quartic divergence and hence the discussion must be postponed until a renormalizable theory of gravity is found.

ACKNOWLEDGMENTS

I thank Professors H.-Y. Guo and S. Chang for the invitation to this Workshop and their kind hospitality extended to me during my stay in Beijing. I have benefitted from discussions with Kiwoon Choi and Hans Peter Nilles. This work is supported in part by Korea Science and Engineering Foundation through Center for Theoretical Physics at Seoul National University, KOSEF-DFG Collaboration Program, and the Ministry of Education of The Republic of Korea through the Basic Science Research Institute, Contract No. BSRI-94-2418.

REFERENCES

- [1] A. Einstein, Sitzungsber. Preuss. Akad. Wiss. Phys.-Math. **K1**, 142 (1917) [English Translation in *Cosmological Constants*, Eds. J. Bernstein and G. Feinberg (Columbia University, New York, 1986)].
- [2] Th. Kaluza, Sitzungsber. Preuss. Akad. Wiss. Berlin, Math. Phys. **K1**, 966 (1921); O. Klein, Z. Phys. **37**, 895 (1926).
- [3] E. Gildener and S. Weinberg, Phys. Rev. **D13**, 3333 (1976).
- [4] K. Choi, J. E. Kim and H. P. Nilles, Phys. Rev. Lett. **73**, 1758 (1994).
- [5] For a review, see, H. P. Nilles, Phys. Rep. **110**, 1 (1984); H. E. Haber and G. Kane, Phys. Rep. **117**, 75 (1985).
- [6] V. Kaplunovski and J. Louis, Phys. Lett. **B306**, 269 (1993); R. Barbieri, J. Louis and M. Moretti, Phys. Lett. **B312**, 451 (1993).
- [7] A. Brignole, L. E. Ibanez and C. Munoz, Univ. Autonoma de Madrid preprint FTUAM-26/93.
- [8] L. Alvarez-Gaume and J. Polchinski, Nucl. Phys. **B221**, 495 (1983).
- [9] A. Manohar and H. Georgi, Nucl. Phys. **B234**, 239 (1984).
- [10] M. Dine and N. Seiberg, Phys. Rev. Lett. **55**, 366 (1985); V. Kaplunovski, Phys. Rev. Lett. **55**, 1036 (1985).
- [11] M. Veltman, Phys. Rev. Lett. **34**, 777 (1975).
- [12] B. Zumino, Nucl. Phys. **B89**, 535 (1975).
- [13] G. F. Giudice and A. Masiero, Phys. Lett. **B206**, 480 (1988).
- [14] See, for example, S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989). For the cosmological constant problem in spontaneously broken gauge theories, see, M. Veltman, Phys. Rev. Lett. **34**, 777 (1975).
- [15] For reviews, see for example, H. P. Nilles, Int. J. Mod. Phys. **A5**, 4199 (1990); D. Amati et al, Phys. Rep. **162**, 169 (1988); J. Louis, in *Proc. 1991 Vancouver DPF Meeting* (World Scientific, Singapore, 1992).
- [16] N. V. Krasnikov, Phys. Lett. **B193**, 37 (1987); L. Dixon, in Proc. 15th APS D.P.F. Meeting, Houston, 1990; J. A. Casas, Z. Lalak, C. Munoz, and G. G. Ross, Nucl. Phys. **B347**, 243 (1990); T. Taylor, Phys. Lett. **B252**, 59 (1990); B. de Carlos, J. A. Casas and C. Munoz, Phys. Lett. **B263**, 248 (1991).
- [17] S. Ferrara, L. Girardello and F. Palumbo, Phys. Rev. **D20**, 403 (1979).
- [18] L. E. Ibanez and D. Lust, Nucl. Phys. **B382**, 305 (1992).
- [19] V. Kaplunovsky and J. Louis, Phys. Lett. **B306**, 269 (1993); R. Barbieri, J. Louis and M. Moretti, Phys. Lett. **B312**, 451 (1993).

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9409420v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9409420v1>

This figure "fig1-3.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9409420v1>